

10581757

File 5:Biosis Previews(R) 1926-2008/Aug W5  
(c) 2008 The Thomson Corporation

Set	Items	Description
---	-----	-----
? s	(distending)	and (campylobacter())coli)
	1209	DISTENDING
	13525	CAMPYLOBACTER
	332256	COLI
	1987	CAMPYLOBACTER(W)COLI
S1	21	(DISTENDING) AND (CAMPYLOBACTER())COLI)

? t s1/7/1-21

1/7/1

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rts. reserv.

0020214933 BIOSIS NO.: 200800261872

Differences in virulence attributes between cytolethal %distending% toxin positive and negative Campylobacter jejuni strains

AUTHOR: Jain Deepika; Prasad Kashi Nath (Reprint); Sinhal Sushmita; Husain Nuzhat

AUTHOR ADDRESS: Sanjay Gandhi Postgrad Inst Med Sci, Dept Microbiol, Lucknow 226014, Uttar Pradesh, India\*\*India

AUTHOR E-MAIL ADDRESS: knprasad@sgpgi.ac.in

JOURNAL: Journal of Medical Microbiology 57 (3): p267-272 MAR 2008 2008

ITEM IDENTIFIER: doi:10.1099/jmm.0.47317-0

ISSN: 0022-2615

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Campylobacter jejuni is a common gastrointestinal bacterial pathogen. Although cytolethal %distending% toxin (CDT) is proposed to be an important virulence determinant of this pathogen, how CDT+ and CDT- strains differ in their biological properties remains largely unknown. The virulence properties of CDT+ and CDT- strains were studied on HeLa cells and in the suckling mouse model. Presence of the cdtB gene in Campylobacter species was determined by PCR. Five each of CDT+ and CDT- C. jejuni strains were subjected to adherence, invasion and cytotoxicity assay on the HeLa cell line. Bacterial culture supernatants with and without CDT activity were inoculated intragastrically into 2-day-old suckling mice. The mice were sacrificed within 48 h. Histopathological examination of stomach, jejunum, ileum and colon was performed by haematoxylin/eosin staining. cdtB was detected in 88 % and 14 % of C. jejuni and %Campylobacter% %coli% strains, respectively. CDT+ C. jejuni strains adhered to and invaded HeLa cells in significantly higher numbers than CDT- strains [CDT+ vs CDT-, adherence  $2.7 \times 10^4$  +/-  $3.5 \times 10^4$  vs  $2.7 \times 10^2$  +/-  $1.9 \times 10^2$ ; invasion  $1.0 \times 10^3$  +/-  $1.3 \times 10^3$  vs  $1.4 \times 10^1$  +/-  $3.1 \times 10^1$ ;  $P < 0.01$ ]. Culture supernatants of all CDT+ strains demonstrated CDT activity on HeLa cells. Mice inoculated with supernatant containing CDT activity had moderate to severe pathology in different parts of their gastrointestinal tract, with the colon being the major target. Mice inoculated with supernatant lacking CDT activity

showed no significant pathology in the gastrointestinal tract. The results demonstrate that CDT+ *C. jejuni* strains adhere to and invade epithelial cells more efficiently than CDT- strains. CDT is responsible for intestinal pathology and the colon is the major target.

1/7/2

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020189327 BIOSIS NO.: 200800236266

Isolation and species identification of campylobacters by genotyping of cytolethal *distending* toxin (CDT) gene from animals

AUTHOR: Asakura M (Reprint); Yoshida E; Samosomsuk W; Sugimoto N; Nishimura K; Yamasaki S

AUTHOR ADDRESS: Univ Osaka Prefecture, Sakai, Osaka 591, Japan\*\*Japan

JOURNAL: Abstracts of the General Meeting of the American Society for Microbiology 105 p141 2005 2005

CONFERENCE/MEETING: 105th General Meeting of the American-Society-for-Microbiology Atlanta, GA, USA June 05 -09, 2005; 20050605

SPONSOR: Amer Soc Microbiol

ISSN: 1060-2011

DOCUMENT TYPE: Meeting; Meeting Abstract

RECORD TYPE: Citation

LANGUAGE: English

1/7/3

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020130234 BIOSIS NO.: 200800177173

Development of a cytolethal *distending* toxin (cdt) gene-based species-specific multiplex PCR assay for the detection and identification of *Campylobacter jejuni*, *Campylobacter coli* and *Campylobacter fetus*

AUTHOR: Asakura Masahiro; Samosornsuk Worada; Hinenoya Atsushi; Misawa Naoaki; Nishimura Kazuhiko; Matsuhisa Akio; Yamasaki Shinji (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Naka Ku, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: FEMS Immunology and Medical Microbiology 52 (2): p260-266 MAR 2008 2008

ITEM IDENTIFIER: doi:10.1111/j.1574-695X.2007.00369.x

ISSN: 0928-8244

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: A cytolethal *distending* toxin (cdt) gene-based species-specific multiplex PCR assay for the detection of *cdtA*, *cdtB* or *cdtC* gene of *Campylobacter jejuni*, *Campylobacter coli* or *Campylobacter fetus*, respectively, was developed and evaluated with 76 *Campylobacter* strains belonging to seven different species and 131 other bacterial strains of eight different genera. The *cdtA*, *cdtB* or *cdtC* gene of *C. jejuni*, *C. coli* or *C. fetus*, respectively, could be successfully

amplified using the corresponding set of primers in a highly species-specific manner. Furthermore, the specific primer set for the *cdtA*, *cdtB* or *cdtC* gene of a particular species could amplify the desired gene from a mixture of DNA templates of any of two or all three species. The detection limit of *C. jejuni*, *C. coli* or *C. fetus* was 10-100 CFU tube(-1) by the multiplex PCR assay on the basis of the presence of the *cdtA*, *cdtB* or *cdtC* gene. These data indicate that the *cdt* gene-based multiplex PCR assay may be useful for rapid and accurate detection as well as identification of *Campylobacter* strains in a species-specific manner.

1/7/4

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020123865 BIOSIS NO.: 200800170804

Optimisation of glycan and small molecule arrays for analysis of *Campylobacter* chemotaxis and adherence

AUTHOR: Asakura M (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Sakai, Osaka, Japan\*\*Japan

JOURNAL: Zoonoses Public Health 54 (Suppl. 1): p99 2007 2007

CONFERENCE/MEETING: 14th International Workshop on *Campylobacter*, *Helicobacter* and Related Organisms Rotterdam, NETHERLANDS September 02-05, 2007; 20070902

ISSN: 1863-1959

DOCUMENT TYPE: Meeting; Meeting Poster

RECORD TYPE: Citation

LANGUAGE: English

1/7/5

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020123846 BIOSIS NO.: 200800170785

Differences in virulence determinants between cytolethal toxin-producing and non-producing *Campylobacter jejuni* strains

AUTHOR: Jain D (Reprint); Prasad K N; Sinha S; Husain N

AUTHOR ADDRESS: SGPGIMS, Lucknow, Uttar Pradesh, India\*\*India

JOURNAL: Zoonoses Public Health 54 (Suppl. 1): p93 2007 2007

CONFERENCE/MEETING: 14th International Workshop on *Campylobacter*, *Helicobacter* and Related Organisms Rotterdam, NETHERLANDS September 02-05, 2007; 20070902

ISSN: 1863-1959

DOCUMENT TYPE: Meeting; Meeting Poster

RECORD TYPE: Citation

LANGUAGE: English

1/7/6

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019895878 BIOSIS NO.: 200700555619

Evaluation of a cytolethal toxin (*cdt*) gene-based species-specific multiplex PCR assay for the identification of

Campylobacter strains isolated from poultry in Thailand  
AUTHOR: Samosornsuk Worada; Asakura Masahiro; Yoshida Emi; Taguchi Takashi;  
Nishimura Kazuhiko; Eampokalap Boonchuay; Phongsisay Vongsavanh;  
Chaicumpa Wanpen; Yamasaki Shinji (Reprint)  
AUTHOR ADDRESS: Osaka Prefecture Univ, Grad Sch Life and Environm Sci, Naka  
Ku, Osaka 5998531, Japan\*\*Japan  
AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp  
JOURNAL: Microbiology and Immunology 51 (9): p909-917 2007 2007  
ISSN: 0385-5600  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: We have recently developed a cytolethal ~~distending~~ toxin (cdt) gene-based species-specific multiplex PCR assay for identifying Campylobacter jejuni, C. coli and C. fetus. In the present study, the applicability of this assay was evaluated with 34 Campylobacter-like organisms isolated from poultry in Thailand for species identification and was compared with other assays including API Campy, 16S rRNA gene sequence, and hippuricase (hipO) gene detection. Of the 34 strains analyzed, 20, 10 and 1 were identified as C. jejuni, C. coli, and Arcobacter cryaerophilus, respectively, and 3 could not be identified by API Campy. However, 16S rRNA gene analysis, showed that all 34 strains are C. jejuni/coli. To discriminate between these 2 species, the hipO gene, which is specifically present in C. jejuni, was examined by PCR and was detected in 20 strains, which were identified as C. jejuni by API Campy but not in the remaining 14 strains. Collective results indicated that 20 strains were C. jejuni whereas the 14 strains were C. coli. When the cdt gene-based multiplex PCR was employed, however, 19, 20 and 19 strains were identified as C. jejuni while 13, 14 and 13 were identified as C. coli by the cdtA, cdtB and cdtC gene-based multiplex PCR, respectively. Pulsed-field gel electrophoresis revealed that C. jejuni and C. coli strains analyzed are genetically diverse. Taken together, these data suggest that the cdt gene-based multiplex PCR, particularly cdtB gene-based multiplex PCR, is a simple, rapid and reliable method for identifying the species of Campylobacter strains.

1/7/7

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rts. reserv.

0019752944 BIOSIS NO.: 200700412685  
Comparative analysis of cytolethal ~~distending~~ toxin (cdt) genes among Campylobacter jejuni, C-coli and C-fetus strains  
AUTHOR: Asakura Masahiro; Samosornsuk Worada; Taguchi Masumi; Kobayashi Kazuhiro; Misawa Naoaki; Kusumoto Masahiro; Nishimura Kazuhiko; Matsuhisa Akio; Yamasaki Shinji (Reprint)  
AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Naka Ku, Gakuen Cho, Sakai, Osaka 5998531, Japan\*\*Japan  
AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp  
JOURNAL: Microbial Pathogenesis 42 (5-6): p174-183 MAY-JUN 2007 2007  
ITEM IDENTIFIER: doi:10.1016/j.micpath.2007.01.005  
ISSN: 0882-4010  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: The cytolethal distending toxin (cdt) gene clusters of *Campylobacter coli* strain Col-243 and *C fetus* strain Col-187 were cloned and sequenced to understand the importance of Cdt as a virulence factor. The cdt genes of *C. coli* and *C fetus* consist of three closely linked genes termed cdtA, cdtB, cdtC whose sizes are 774, 801, and 570 bp, and 702, 798, and 546 bp, respectively. The homologies of each subunit of cdt genes between *C jejuni* and *C coli*, *C jejuni* and *C fetus*, or *C coli* and *C fetus* are 59.6%, 40.3%, or 46.5% for cdtA, 70.2%, 62.4%, or 61.3% for cdtB, 61.3%, 52.3%, or 50.1% for cdtC, respectively. Colony hybridization assay revealed that the genes homologous to the cdtABC gene were distributed in all 27, 19, 20 strains of *C jejuni*, *C. coli*, and *C fetus*, respectively, isolated from patients and animals in species-specific manner. Furthermore, nucleotide sequence of the cdt operon, including flanking region, of 10 strains of each species indicated that though the size of the cdtB gene was conserved in each species, those of cdtA and cdtC genes varied particularly among *C coli* strains. Amino acid residues demonstrated to be important for toxin activity in CdtB, corresponding to H 152, D185, D222, D258, H259 in Cj-CdtB, were also conserved in Cc-CdtB and Cf-CdtB. The cdt gene cluster was located in different sites among different species but in the same site of genomes of the same species. Cdt activity produced by *C jejuni* and *C. fetus* varied among strains, however, any *C coli* strains exhibited Cdt activity on HeLa cells. These data indicate that the cdt gene may have a potential for virulence factor at least in *C jejuni* and *C fetus*.  
(C) 2007 Elsevier Ltd. All rights reserved.

1/7/8

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

0019664241 BIOSIS NO.: 200700323982

Development of a multiplex PCR assay for the detection of the cytolethal distending toxin genes in *Campylobacter jejuni*, *C-coli* and *C-fetus*

AUTHOR: Asakura M (Reprint); Yoshida E; Nishimura K; Taguchi A; Kobayashi K; Yamasaki S

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Agr and Biol Sci, Osaka, Japan\*\*Japan

JOURNAL: Abstracts of the General Meeting of the American Society for Microbiology 104 p209 2004 2004

CONFERENCE/MEETING: 104th General Meeting of the American-Society-for-Microbiology New Orleans, LA, USA May 23 -27, 2004; 20040523

SPONSOR: Amer Soc Microbiol

ISSN: 1060-2011

DOCUMENT TYPE: Meeting; Meeting Abstract

RECORD TYPE: Citation

LANGUAGE: English

1/7/9

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

0019544568 BIOSIS NO.: 200700204309

Relationships between bacterial genotypes and in vitro virulence properties

of *Campylobacter jejuni* and *Campylobacter coli* isolated from turkeys

AUTHOR: Haenel I (Reprint); Borrmann E; Mueller J; Alter T

AUTHOR ADDRESS: Fed Res Inst Anim Hlth, Inst Mol Pathogenesis, Naumburger Str 96A, D-07743 Jena, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: ingrid.haenel@fli.bund.de

JOURNAL: Journal of Applied Microbiology 102 (2): p433-441 FEB 2007 2007

ISSN: 1364-5072

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: *Campylobacter* isolates from turkeys were genotyped and characterized by their in vitro virulence properties. Relationships between bacterial genotypes and virulence properties were analysed. Isolates were analysed by pulsed-field gel electrophoresis and fla typing. The toxin production was determined on the phenotypic level using a CHO-K1 cell culture model and on the genotypic level using PCR for detection of the *cdtA*, *cdtB* and *cdtC* genes. Although the *cdtB* gene was detected from 100% of the *Campylobacter jejuni* and *Campylobacter coli* isolates we observed three different morphological pictures on the cells. Cytotoxicity was associated with cell distension or cell rounding. All four *Camp. coli* strains and one *Camp. jejuni* strain did not produce any cytotoxic changes on the cells. Adhesion, invasion and survival of *Campylobacter* isolates were determined in a Caco-2 cell culture model. All isolates adhered to and invaded Caco-2 cells, whereas 64.7% of the strains survived for 48 h in the cells. Seventeen *Campylobacter* isolates from turkeys were classified into four groups with regard to their in vitro abilities. Jackknife analysis revealed a strong association between these groups and genotype clusters. Typing methods have generally failed to identify strains with specific virulence properties. This study suggests that a relationship between subgroups of *Campylobacter* with common in vitro virulence characteristics and genotypes exist.

1/7/10

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

19060686 BIOSIS NO.: 200600406081

Adherence to and invasion of human intestinal epithelial cells by

*Campylobacter jejuni* and *Campylobacter coli* isolates from retail meat products

AUTHOR: Zheng JIE; Meng JIANGHONG; Zhao SHAOHUA; Singh RUBY; Song WENXIA (Reprint)

AUTHOR ADDRESS: Univ Maryland, Dept Cell Biol and Mol Genet, College Pk, MD 20742 USA\*\*USA

AUTHOR E-MAIL ADDRESS: wenxsong@umd.edu

JOURNAL: Journal of Food Protection 69 (4): p768-774 APR 2006 2006

ISSN: 0362-028X

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The abilities of 34 *Campylobacter jejuni* and 9

*Campylobacter coli* isolates recovered from retail meats to

adhere to and invade human intestinal epithelial T84 cells were examined and compared with those of a well-characterized human clinical strain, C jejuni 81-176, to better assess the pathogenic potential of these meat isolates. The meat isolates exhibited a wide range of adherence and invasion abilities; a few of the isolates adhered to and invaded T84 cells almost as well as did C jejuni 81-176. There was a significant correlation between the adherence ability and the invasion ability of the Campylobacter isolates. The presence of eight putative virulence genes in these Campylobacter isolates that are potentially responsible for adherence and invasion or that encode cytolethal distending toxin was determined using PCR. All Campylobacter isolates possessed flaA, cadF, pldA, cdtA, cdtB, and cdtC, and most (91%) also contained the ciaB gene. However, the virBII gene, carried by virulence plasmid pVir, was absent in almost all the Campylobacter isolates. Our findings indicated that C jejuni and C. coli present in retail meat were diverse in their ability to adhere to and invade human intestinal epithelial cells and that the putative virulence genes were widespread among the Campylobacter isolates. Thus, despite of the presence of the putative virulence genes, only some but not all Campylobacter strains isolated from retail meat can effectively invade human intestinal epithelial cells in vitro.

1/7/11

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

17391700 BIOSIS NO.: 200300350419

PCR detection of seven virulence and toxin genes of Campylobacter jejuni and Campylobacter coli isolates from Danish pigs and cattle and cytolethal distending toxin production of the isolates.

AUTHOR: Bang D D (Reprint); Nielsen E Moller; Scheutz F; Pedersen K; Handberg K; Madsen M

AUTHOR ADDRESS: Department of Poultry, Fish and Fur Animals, Danish Veterinary Institute, Hangevej 2, DK-8200, Aarhus N, Denmark\*\*Denmark

AUTHOR E-MAIL ADDRESS: ddb@vetinst.dk

JOURNAL: Journal of Applied Microbiology 94 (6): p1003-1014 2003 2003

MEDIUM: print

ISSN: 1364-5072

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Aims: To study the prevalence of seven virulence and toxin genes, and cytolethal distending toxin (CDT) production of Campylobacter jejuni and C. coli isolates from Danish pigs and cattle. Methods and Results: The presence of the cadF, ceuE, virB11, flaA, cdtA, cdtB, cdtC and the cdt gene cluster among 40 C. jejuni and C. coli isolates was detected by polymerase chain reaction. The CDT production of the isolates was determined on Vero, colon 205 and chicken embryo cells. The cadF, flaA, ceuE and cdtB genes were detected from 100% of the isolates. The cdtA and cdtC genes were found in 95.0 and 90.0% of the isolates, respectively. The cdt gene cluster was detected in 82.5% isolates. Only 7.5% of the isolates were positive for virB11. Ninety-five per cent of the isolates produced CDT in Vero and colon 205 cell assays, and 90% of the isolates produced CDT in chicken embryo cell assays. Conclusions: High prevalence of the cadF, ceuE, flaA and cdtB genes was found. Data of the prevalence of cdt genes was consistent with the CDT titres produced

by the isolates. *Campylobacter coli* from pigs produced high CDT titres. Significance and Impact of the Study: The high prevalence of seven virulence and toxin genes demonstrated that these putative pathogenic determinants are widespread among *Campylobacter* isolates from pigs and cattle. *Campylobacter coli* isolates from pigs produced much higher CDT titres compared with *C. coli* isolates from other sources suggesting that *C. coli* may be particularly adapted to or associated with this species.

1/7/12

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

16846487 BIOSIS NO.: 200200439998

*Campylobacter jejuni* cytolethal *distending* toxin mediates release of interleukin-8 from intestinal epithelial cells

AUTHOR: Hickey Thomas E; McVeigh Annette L; Scott Daniel A; Michielutti Ronda E; Bixby Alyssa; Carroll Shannon A; Bourgeois A Louis; Guerry Patricia (Reprint)

AUTHOR ADDRESS: Enteric Diseases Department, Naval Medical Research Center, 503 Robert Grant Ave., Walter Reed Forest Glen Annex, Silver Spring, MD, 20910, USA\*\*USA

JOURNAL: Infection and Immunity 68 (12): p6535-6541 December, 2000 2000

MEDIUM: print

ISSN: 0019-9567

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Live cells of *Campylobacter jejuni* and *Campylobacter coli* can induce release of interleukin-8 (IL-8) from INT407 cells. Additionally, membrane fractions of *C. jejuni* 81-176, but not membrane fractions of *C. coli* strains, can also induce release of IL-8. Membrane preparations from 81-176 mutants defective in any of the three membrane-associated protein subunits of cytolethal *distending* toxin (CDT) were unable to induce IL-8. The presence of the three *cdt* genes on a shuttle plasmid in trans restored both CDT activity and the ability to release IL-8 to membrane fractions. However, CDT mutations did not affect the ability of 81-176 to induce IL-8 during adherence to or invasion of INT407 cells. When *C. jejuni* *cdt* genes were transferred on a shuttle plasmid into a *C. coli* strain lacking CDT, membrane preparations became positive in both CDT and IL-8 assays. Growth of *C. jejuni* in physiological levels of sodium deoxycholate released all three CDT proteins, as well as CDT activity and IL-8 activity, from membranes into supernatants. Antibodies against recombinant forms of each of the three CDT subunit proteins neutralized both CDT activity and the activity responsible for IL-8 release. The data suggest that *C. jejuni* can induce IL-8 release from INT407 cells by two independent mechanisms, one of which requires adherence and/or invasion and the second of which requires CDT.

1/7/13

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.



16479692 BIOSIS NO.: 200200073203

Prevalence of cytolethal distending toxin (cdt) genes and CDT production in *Campylobacter* spp. isolated from Danish broilers

AUTHOR: Bang Dang D (Reprint); Scheutz Flemming; Ahrens Peter; Pedersen Karl; Blom Jens; Madsen Mogens

AUTHOR ADDRESS: Department of Poultry, Fish, and Fur Animals, Danish Veterinary Laboratory, Hangovej 2, DK-8200, Aarhus N, Denmark\*\*Denmark

JOURNAL: Journal of Medical Microbiology 50 (12): p1087-1094 December, 2001 2001

MEDIUM: print

ISSN: 0022-2615

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The pathogenesis of campylobacter infection in man is largely unknown, although cytolethal distending toxin (CDT) has been incriminated as a virulence factor. However, little is known about the cdt genes in *Campylobacter* spp. isolated from broiler chickens. A total of 350 cloacal swabs was collected and tested by conventional culture and PCR. Of the 114 *Campylobacter* isolates obtained, 101 (88.6%) were identified as *C. jejuni* and 13 (11.4%) as *C. coli* by conventional methods. cdt genes were detected by PCR in all the isolates except one *C. jejuni* isolate. Cytotoxic effects were produced in a Vero cell line, by 100 of the *C. jejuni* isolates. In contrast, 10 *C. coli* isolates produced much lower levels of toxin and 3 produced no detectable toxin. These results confirm the common occurrence of campylobacter infection in chickens and indicate that cdt genes are commonly present in both *C. jejuni* and *C. coli* isolates from broilers, but that there are distinct differences in CDT production in these two closely related species.

1/7/14

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

16477619 BIOSIS NO.: 200200071130

Classical and molecular identification of *Campylobacter jejuni* and *Campylobacter coli* from poultry samples on Slovenian market

AUTHOR: Zorman T (Reprint); Mavri U; Mozina S Smole

AUTHOR ADDRESS: Biotechnical Faculty, Department for Food Science and Technology, Slovenia, University of Ljubljana, Ljubljana, Slovenia\*\*Slovenia

JOURNAL: IJMM International Journal of Medical Microbiology 291 (Supplement 31): p40 September, 2001 2001

MEDIUM: print

CONFERENCE/MEETING: 11th International Workshop on *Campylobacter*, *Helicobacter* and related Organisms Freiburg, Germany September 01-05, 2001; 20010901

ISSN: 1438-4221

DOCUMENT TYPE: Meeting; Meeting Abstract

RECORD TYPE: Citation

LANGUAGE: English

1/7/15

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

16460440 BIOSIS NO.: 200200053951  
Cytolethal %distending% toxin B gene (cdtB) homologues in taxons 1, 2,  
3, 4, 5 and 8 of Helicobacter species flexispira  
AUTHOR: Kostia S (Reprint); Hanninen M (Reprint)  
AUTHOR ADDRESS: University of Helsinki, Helsinki, Finland\*\*Finland  
JOURNAL: IJMM International Journal of Medical Microbiology 291 (  
Supplement 31): p149 September, 2001 2001  
MEDIUM: print  
CONFERENCE/MEETING: 11th International Workshop on Campylobacter,  
Helicobacter and related Organisms Freiburg, Germany September 01-05,  
2001; 20010901  
ISSN: 1438-4221  
DOCUMENT TYPE: Meeting; Meeting Abstract  
RECORD TYPE: Citation  
LANGUAGE: English

1/7/16

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rts. reserv.

15168000 BIOSIS NO.: 199900427660  
The cytolethal %distending% toxin family  
AUTHOR: Pickett Carol L (Reprint); Whitehouse Chris A (Reprint)  
AUTHOR ADDRESS: Dept of Microbiology and Immunology, University of  
Kentucky, Lexington, KY, 40536-0298, USA\*\*USA  
JOURNAL: Trends in Microbiology 7 (7): p292-297 July, 1999 1999  
MEDIUM: print  
ISSN: 0966-842X  
DOCUMENT TYPE: Article; Literature Review  
RECORD TYPE: Citation  
LANGUAGE: English

1/7/17

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rts. reserv.

14984828 BIOSIS NO.: 199900244488  
Detection of cytolethal %distending% toxin activity and cdt genes in  
Campylobacter spp. isolated from chicken carcasses  
AUTHOR: Eyigor Aysegul; Dawson Karl A; Langlois Bruce E; Pickett Carol L  
(Reprint)  
AUTHOR ADDRESS: Department of Microbiology and Immunology, Chandler Medical  
Center, University of Kentucky, 800 Rose St., Lexington, KY, 40536-0084,  
USA\*\*USA  
JOURNAL: Applied and Environmental Microbiology 65 (4): p1501-1505 April,  
1999 1999  
MEDIUM: print  
ISSN: 0099-2240  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: This study was designed to determine whether isolates from

chicken carcasses, the primary source of *Campylobacter jejuni* and *Campylobacter coli* in human infections, commonly carry the *cdt* genes and also whether active cytolethal distending toxin (CDT) is produced by these isolates. *Campylobacter* spp. were isolated from all 91 fresh chicken carcasses purchased from local supermarkets. *Campylobacter* spp. were identified on the basis of both biochemical and PCR tests. Of the 105 isolates, 70 (67%) were identified as *C. jejuni*, and 35 (33%) were identified as *C. coli*. PCR tests amplified portions of the *cdt* genes from all 105 isolates. Restriction analysis of PCR products indicated that there appeared to be species-specific differences between the *C. jejuni* and *C. coli* *cdt* genes, but that the restriction patterns of the *cdt* genes within strains of the same species were almost invariant. Quantitation of active CDT levels produced by the isolates indicated that all *C. jejuni* strains except four (94%) had mean CDT titers greater than 100. Only one *C. jejuni* strain appeared to produce no active CDT. *C. coli* isolates produced little or no toxin. These results confirm the high rate of *Campylobacter* sp. contamination of fresh chicken carcasses and indicate that *cdt* genes may be universally present in *C. jejuni* and *C. coli* isolates from chicken carcasses.

1/7/18

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

14977308 BIOSIS NO.: 199900236968

Cytolethal distending toxin genes in *Campylobacter jejuni* and *Campylobacter coli* isolates: Detection and analysis by PCR

AUTHOR: Eyigor Aysegul; Dawson Karl A; Langlois Bruce E; Pickett Carol L (Reprint)

AUTHOR ADDRESS: Department of Microbiology and Immunology, Chandler Medical Center, University of Kentucky, 800 Rose St., Lexington, KY, 40536-0298, USA\*\*USA

JOURNAL: Journal of Clinical Microbiology 37 (5): p1646-1650 May, 1999

MEDIUM: print

ISSN: 0095-1137

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: *Campylobacter jejuni* produces a toxin called cytolethal distending toxin (CDT). Knowledge of the prevalence and homogeneity of *Campylobacter* sp. *cdt* genes is incomplete. In this work, we identified four PCR primer pairs that collectively amplified *cdt* genes in all of the *C. jejuni* and *Campylobacter coli* strains tested. Restriction analyses of the *cdt* PCR products showed clear differences between the *cdt* genes of these two species, yet there were few heterogeneities noted between members of the same species. Consequently, it may be possible to speciate *C. jejuni* and *C. coli* isolates on the basis of restriction patterns within their *cdt* genes.

1/7/19

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

14623418 BIOSIS NO.: 199800417665

Campylobacter spp. isolated from chicken carcasses: Prevalence and detection of cytolethal ~~distending~~ toxin production

AUTHOR: Eyigor A; Langlois B E; Dawson K; Pickett C L

AUTHOR ADDRESS: Univ. Kentucky, Lexington, KY, USA\*\*USA

JOURNAL: Abstracts of the General Meeting of the American Society for Microbiology 98 p413 1998 1998

MEDIUM: print

CONFERENCE/MEETING: 98th General Meeting of the American Society for Microbiology Atlanta, Georgia, USA May 17-21, 1998; 19980517

SPONSOR: American Society for Microbiology

ISSN: 1060-2011

DOCUMENT TYPE: Meeting; Meeting Abstract; Meeting Poster

RECORD TYPE: Citation

LANGUAGE: English

1/7/20

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

13406673 BIOSIS NO.: 199699040733

Prevalence of cytolethal ~~distending~~ toxin production in Campylobacter jejuni and relatedness of Campylobacter sp. cdtB genes

AUTHOR: Pickett Carol L (Reprint); Pesci Evertt C; Cottle Daniel L; Russell Gina; Erdem Aysegul Nalca; Zeytn Hasan

AUTHOR ADDRESS: Dep. Microbiol. Immunol., Univ. Kentucky, Chandler Med. Cent., 800 Rose Street, Lexington, KY 40536-0084, USA\*\*USA

JOURNAL: Infection and Immunity 64 (6): p2070-2078 1996 1996

ISSN: 0019-9567

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Campylobacter jejuni produces a toxin called cytolethal ~~distending~~ toxin (CDT). The genes encoding this toxin in C. jejuni 81-176 were cloned and sequenced. The nucleotide sequence of the genes revealed that there are three genes, cdtA, cdtB, and cdtC, encoding proteins with predicted sizes of 30,116, 28,989, and 21,157 Da, respectively. All three proteins were found to be related to the Escherichia coli CDT proteins, yet the amino acid sequences have diverged significantly. All three genes were required for toxic activity in a HeLa cell assay. HeLa cell assays of a variety of C. jejuni and C. coli strains suggested that most C. jejuni strains produce significantly higher CDT titers than do C. coli strains. Southern hybridization experiments demonstrated that the cdtB gene is present on a 6.0-kb ClaI fragment in all but one of the C. jejuni strains tested; the cdtB gene was on a 6.9-kb ClaI fragment in one strain. The C. jejuni 81-176 cdtB probe hybridized weakly to DNAs from C. coli strains. The C. jejuni 81-176 cdtB probe did not hybridize to DNAs from representative C. fetus, C. lari, C. "upsaliensis," and C. hyointestinalis strains, although the HeLa cell assay indicated that these strains make CDT. PCR experiments indicated the probable presence of cdtB sequences in all of these Campylobacter species.

1/7/21

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rts. reserv.

09176631 BIOSIS NO.: 198886016552

A NEW HEAT-LABILE CYTOLETHAL %%%DISTENDING%%% TOXIN CLDT PRODUCED BY  
CAMPYLOBACTER-SPP

AUTHOR: JOHNSON W M (Reprint); LIOR H

AUTHOR ADDRESS: NATL ENTERIC REFERENCE CENT, ENTERIC BACTERIOL DIV, BUREAU  
MICROBIOLOGY, LAB CENTRE DIS CONTROL, TUNNEY'S PASTURE, OTTAWA, ONT K1A  
0L2, CAN\*\*CANADA

JOURNAL: Microbial Pathogenesis 4 (2): p115-126 1988

ISSN: 0882-4010

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: ENGLISH

ABSTRACT: A new heat-labile toxin cytolethal to CHO, Vero, HeLa, and HEp-2 cells and negative in Y-1 cells has been demonstrated in culture filtrates of many strains of Campylobacter spp. This new toxin was termed a cytolethal %%%distending%%% toxin (CLDT) to reflect the progressive cell distention and eventual cytotoxicity observed in all sensitive tissue cells. CLDT was distinct from previously reported cytotoxins and cholera-like enterotoxin produced by some Campylobacter spp. Since CHO elongation induced by either the Campylobacter enterotoxin or CLDT could not be differentiated after 2 h incubation, continuation of the assay for 96 h was essential for observation of CLDT-associated progressive morphological changes and cytolethal events. Specific assay conditions were required for demonstration of CLDT in Vero, HeLa, and HEp-2 cells. A 31-fold increase in cyclic AMP levels was observed in CHO cells exposed for 24 h to CLDT of catalase negative or weak positive Campylobacter. CLDT was detected in culture filtrates from strains of Campylobacter jejuni, c. coli, C. fetus subsp. fetus, C. laridis and catalase negative or weak positive Campylobacter. Of 718 strains investigated from both human and animal isolations, 295 (41%) were found to produce this toxin. Campylobacter CLDT was negative in adult rabbit ligated ileal loops, suckling mouse and rabbit skin tests. Hemorrhagic responses were observed in rat ligated ileal loop tests of CLDT-positive cultures. The new CLDT toxin could only be neutralized by homologous rabbit antitoxin, was trypsin-sensitive, nondialyzable and over 30,000 in molecular weight. CLDT-producing strains were observed in many serogroups and biotypes of Campylobacter spp. The strains tested originated in many countries and no clear association of toxigenicity with serotype or biotype could be established.

? ds

Set	Items	Description
S1	21	(DISTENDING) AND (CAMPYLOBACTER())COLI)

? e au=yamasaki shinji

Ref	Items	Index-term
E1	3	AU=YAMASAKI SHINICHI
E2	1	AU=YAMASAKI SHINICHIROU
E3	87	*AU=YAMASAKI SHINJI
E4	1	AU=YAMASAKI SHINSUKE
E5	9	AU=YAMASAKI SHINTARO
E6	1	AU=YAMASAKI SHINZO
E7	2	AU=YAMASAKI SHIORI

E8 4 AU=YAMASAKI SHIROU  
 E9 35 AU=YAMASAKI SHO  
 E10 1 AU=YAMASAKI SHOGO  
 E11 4 AU=YAMASAKI SHOICHI  
 E12 1 AU=YAMASAKI SHOUSAKU

Enter P or PAGE for more

? s e3

S2 87 AU='YAMASAKI SHINJI'

? e au=assakura masahiro

Ref	Items	Index-term
E1	11	AU=ASSAKURA MARINA T
E2	1	AU=ASSAKURA MARINA TIZUKO
E3	0	*AU=ASSAKURA MASAHIRO
E4	13	AU=ASSAL A
E5	3	AU=ASSAL A A
E6	1	AU=ASSAL A M
E7	2	AU=ASSAL A N
E8	3	AU=ASSAL AMY A
E9	2	AU=ASSAL AZZEDINE
E10	2	AU=ASSAL C
E11	1	AU=ASSAL E
E12	11	AU=ASSAL F

Enter P or PAGE for more

? e au= asakura masahiro

Ref	Items	Index-term
E1	1	AU=ASAKURA MAMORU
E2	3	AU=ASAKURA MASAE
E3	15	*AU=ASAKURA MASAHIRO
E4	92	AU=ASAKURA MASANORI
E5	3	AU=ASAKURA MASATO
E6	3	AU=ASAKURA MASUMI
E7	1	AU=ASAKURA MIKA
E8	1	AU=ASAKURA MIKI
E9	15	AU=ASAKURA MIKIO
E10	1	AU=ASAKURA MITSUHIITO
E11	14	AU=ASAKURA N
E12	1	AU=ASAKURA NAOKO

Enter P or PAGE for more

? s e3

S3 15 AU='ASAKURA MASAHIRO'

? s s2 and s3

87 S2

15 S3

S4 10 S2 AND S3

? t s4/7/1-10

4/7/1

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020130234 BIOSIS NO.: 200800177173

Development of a cytolethal distending toxin (cdt) gene-based

species-specific multiplex PCR assay for the detection and identification of *Campylobacter jejuni*, *Campylobacter coli* and *Campylobacter fetus*  
AUTHOR: %Asakura Masahiro%; Samosornsuk Worada; Hinenoya Atsushi; Misawa Naoaki; Nishimura Kazuhiko; Matsuhisa Akio; %Yamasaki Shinji% (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Naka Ku, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: FEMS Immunology and Medical Microbiology 52 (2): p260-266 MAR 2008 2008

ITEM IDENTIFIER: doi:10.1111/j.1574-695X.2007.00369.x

ISSN: 0928-8244

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: A cytolethal distending toxin (cdt) gene-based species-specific multiplex PCR assay for the detection of *cdtA*, *cdtB* or *cdtC* gene of *Campylobacter jejuni*, *Campylobacter coli* or *Campylobacter fetus*, respectively, was developed and evaluated with 76 *Campylobacter* strains belonging to seven different species and 131 other bacterial strains of eight different genera. The *cdtA*, *cdtB* or *cdtC* gene of *C. jejuni*, *C. coli* or *C. fetus*, respectively, could be successfully amplified using the corresponding set of primers in a highly species-specific manner. Furthermore, the specific primer set for the *cdtA*, *cdtB* or *cdtC* gene of a particular species could amplify the desired gene from a mixture of DNA templates of any of two or all three species. The detection limit of *C. jejuni*, *C. coli* or *C. fetus* was 10-100 CFU tube(-1) by the multiplex PCR assay on the basis of the presence of the *cdtA*, *cdtB* or *cdtC* gene. These data indicate that the *cdt* gene-based multiplex PCR assay may be useful for rapid and accurate detection as well as identification of *Campylobacter* strains in a species-specific manner.

4/7/2

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019923467 BIOSIS NO.: 200700583208

An inducible lambdoid prophage encoding cytolethal distending toxin (Cdt-I) and a type III effector protein in enteropathogenic *Escherichia coli*

AUTHOR: %Asakura Masahiro%; Hinenoya Atsushi; Alam Mohammad S; Shima Kensuke; Zahid Shamim Hasan; Shi Lei; Sugimoto Norihiko; Ghosh A N; Ramamurthy T; Faruque Shah M; Nair G Balakrish (Reprint); %Yamasaki% % Shinji%

AUTHOR ADDRESS: Int Ctr Diarrhoeal Dis Res, Mol Genet Lab, Dhaka 1212, Bangladesh\*\*Bangladesh

AUTHOR E-MAIL ADDRESS: gbnair@icddr.org; shinji@vet.osakafu-u.ac.jp

JOURNAL: Proceedings of the National Academy of Sciences of the United States of America 104 (36): p14483-14488 SEP 4 2007 2007

ITEM IDENTIFIER: doi:10.1073/pnas.0706695104

ISSN: 0027-8424

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Cytolethal distending toxins (CDTs) are inhibitory cyclomodulins,

which block eukaryotic cell proliferation and are produced by a diverse group of Gram-negative bacteria, including *Escherichia coli* strains associated with intestinal and extraintestinal infections. However, the mode of transmission of the toxin gene clusters among diverse bacterial pathogens is unclear. We found that Cdt-I produced by enteropathogenic *E. coli* strains associated with diarrhea is encoded by a lambdoid prophage, which is inducible and infectious. The genome of Cdt-I converting phage (CDT-14) comprises 47,021 nucleotides with 60 predicted ORFs organized into six genomic regions encoding the head and tail, virulence, integrase, unknown functions, regulation, and lysis. The genomic organization of CDT-1(D is similar to those of SfV, a serotype-converting phage of *Shigella flexneri*, and UT189, a prophage identified in uropathogenic *E. coli*. Besides the *cdtI* gene cluster, the virulence region of CDT-1(P genome contains sequences homologous to a truncated cycle inhibiting factor and a type 3 effector protein. Mutation analysis of susceptible *E. coli* strain C600 suggested that the outer membrane protein OmpC is a putative receptor for CDT-1(D. CDT-1 Phi genome was also found to integrate into the host bacterial chromosome forming lysogens, which produced biologically active Cdt-1. Furthermore, phage induction appeared to cause enhanced toxigenicity of the *E. coli* strains carrying lysogenic CDT-1(D. Our results suggest that CDT-14) is the latest member of a growing family of lambdoid phages encoding bacterial cyclomodulins and that the phage may have a role in horizontal transfer of these virulence genes.

4/7/3

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019895878 BIOSIS NO.: 200700555619

Evaluation of a cytolethal distending toxin (*cdt*) gene-based species-specific multiplex PCR assay for the identification of *Campylobacter* strains isolated from poultry in Thailand

AUTHOR: Samosornsuk Worada; %%Asakura Masahiro%%; Yoshida Emi; Taguchi Takashi; Nishimura Kazuhiko; Eampokalap Boonchuay; Phongsisay Vongsavanh; Chaicumpa Wanpen; %%Yamasaki Shinji%% (Reprint)

AUTHOR ADDRESS: Osaka Prefecture Univ, Grad Sch Life and Environm Sci, Naka Ku, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbiology and Immunology 51 (9): p909-917 2007 2007

ISSN: 0385-5600

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: We have recently developed a cytolethal distending toxin (*cdt*) gene-based species-specific multiplex PCR assay for identifying *Campylobacter jejuni*, *C. coli* and *C. fetus*. In the present study, the applicability of this assay was evaluated with 34 *Campylobacter*-like organisms isolated from poultry in Thailand for species identification and was compared with other assays including API Campy, 16S rRNA gene sequence, and hippuricase (*hipO*) gene detection. Of the 34 strains analyzed, 20, 10 and 1 were identified as *C. jejuni*, *C. coli*, and *Arcobacter cryaerophilus*, respectively, and 3 could not be identified by API Campy. However, 16S rRNA gene analysis, showed that all 34 strains are *C. jejuni/coli*. To discriminate between these 2 species, the *hipO*



gene, which is specifically present in *C. jejuni*, was examined by PCR and was detected in 20 strains, which were identified as *C. jejuni* by API Campy but not in the remaining 14 strains. Collective results indicated that 20 strains were *C. jejuni* whereas the 14 strains were *C. coli*. When the *cdt* gene-based multiplex PCR was employed, however, 19, 20 and 19 strains were identified as *C. jejuni* while 13, 14 and 13 were identified as *C. coli* by the *cdtA*, *cdtB* and *cdtC* gene-based multiplex PCR, respectively. Pulsed-field gel electrophoresis revealed that *C. jejuni* and *C. coli* strains analyzed are genetically diverse. Taken together, these data suggest that the *cdt* gene-based multiplex PCR, particularly *cdtB* gene-based multiplex PCR, is a simple, rapid and reliable method for identifying the species of *Campylobacter* strains.

4/7/4

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019752944 BIOSIS NO.: 200700412685

Comparative analysis of cytolethal distending toxin (*cdt*) genes among *Campylobacter jejuni*, *C. coli* and *C. fetus* strains

AUTHOR: %%Asakura Masahiro%%; Samosornsuk Worada; Taguchi Masumi; Kobayashi Kazuhiro; Misawa Naoaki; Kusumoto Masahiro; Nishimura Kazuhiko; Matsuhisa Akio; %%Yamasaki Shinji%% (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Naka Ku, Gakuen Cho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbial Pathogenesis 42 (5-6): p174-183 MAY-JUN 2007 2007

ITEM IDENTIFIER: doi:10.1016/j.micpath.2007.01.005

ISSN: 0882-4010

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The cytolethal distending toxin (*cdt*) gene clusters of *Campylobacter coli* strain Col-243 and *C. fetus* strain Col-187 were cloned and sequenced to understand the importance of Cdt as a virulence factor. The *cdt* genes of *C. coli* and *C. fetus* consist of three closely linked genes termed *cdtA*, *cdtB*, *cdtC* whose sizes are 774, 801, and 570 bp, and 702, 798, and 546 bp, respectively. The homologies of each subunit of *cdt* genes between *C. jejuni* and *C. coli*, *C. jejuni* and *C. fetus*, or *C. coli* and *C. fetus* are 59.6%, 40.3%, or 46.5% for *cdtA*, 70.2%, 62.4%, or 61.3% for *cdtB*, 61.3%, 52.3%, or 50.1% for *cdtC*, respectively. Colony hybridization assay revealed that the genes homologous to the *cdtABC* gene were distributed in all 27, 19, 20 strains of *C. jejuni*, *C. coli*, and *C. fetus*, respectively, isolated from patients and animals in species-specific manner. Furthermore, nucleotide sequence of the *cdt* operon, including flanking region, of 10 strains of each species indicated that though the size of the *cdtB* gene was conserved in each species, those of *cdtA* and *cdtC* genes varied particularly among *C. coli* strains. Amino acid residues demonstrated to be important for toxin activity in CdtB, corresponding to H152, D185, D222, D258, H259 in Cj-CdtB, were also conserved in Cc-CdtB and Cf-CdtB. The *cdt* gene cluster was located in different sites among different species but in the same site of genomes of the same species. Cdt activity produced by *C. jejuni* and *C. fetus* varied among strains, however, any *C. coli* strains exhibited Cdt activity on HeLa cells. These data indicate that the *cdt* gene may

have a potential for virulence factor at least in *C jejuni* and *C fetus*.  
(C) 2007 Elsevier Ltd. All rights reserved.

4/7/5

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019685440 BIOSIS NO.: 200700345181

Cytotolethal distending toxin (Cdt)-producing *Escherichia coli* isolated from  
a child with bloody diarrhea in Japan

AUTHOR: Hinenoya Atsushi; Nagita Akira; %%Asakura Masahiro%%; Tsukamoto  
Teizo; Ramamuthy Thandavarayan; Nair Gopinath Balakrish; Takeda Yoshifumi  
; %%Yamasaki Shinji%% (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci,  
Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbiology and Immunology 51 (4): p435-438 2007 2007

ISSN: 0385-5600

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: In a retrospective analysis by PCR, the *cdtI* gene encoding the  
cytotolethal distending toxin (Cdt) was detected in *Escherichia coli* O2:H12  
strain isolated from the bloody diarrheal stool specimen of a child. To  
our knowledge, this is the first report showing the possible association  
of Cdt-producing *E. coli* in Japan, particularly in a child with bloody  
diarrhea.

4/7/6

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

19097665 BIOSIS NO.: 200600443060

Species-specific identification of *Vibrio fluvialis* by PCR targeted to the  
conserved transcriptional activation and variable membrane tether regions  
of the *toxR* gene

AUTHOR: Chakraborty Rupa; Sinha Sutapa; Mukhopadhyay Asish K; %%Asakura%%  
%%Masahiro%%; %%Yamasaki Shinji%%; Bhattacharya S K; Nair G Balakrish;  
Ramamurthy T (Reprint)

AUTHOR ADDRESS: Natl Inst Cholera and Enter Dis, P-33,CIT Rd,Scheme XM,  
Calcutta 700010, India\*\*India

AUTHOR E-MAIL ADDRESS: tramu@vsnl.net

JOURNAL: Journal of Medical Microbiology 55 (6): p805-808 JUN 2006 2006

ISSN: 0022-2615

DOCUMENT TYPE: Letter; Editorial

RECORD TYPE: Citation

LANGUAGE: English

4/7/7

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

19087243 BIOSIS NO.: 200600432638

Effects of polyamines on two strains of *Trypanosoma brucei* in infected rats and in vitro culture

AUTHOR: Nishimura Kazuhiko (Reprint); Yanase Takako; Araki Noriko; Ohnishi Yoshihiro; Kozaki Shunji; Shima Kensuke; %Asakura Masahiro%; Samosomsuk Worada; %Yamasaki Shinji%

AUTHOR ADDRESS: Univ Osaka Prefecture, Course Vet Sci, Grad Sch Life and Environm Sci, 1-1, Gakuencho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: nisimura@vet.osakafu-u.ac.jp

JOURNAL: Journal of Parasitology 92 (2): p211-217 APR 2006 2006

ISSN: 0022-3395

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: We studied the effects of polyamines, which are necessary for proliferation and antioxidation in *Trypanosoma brucei gambiense* Wellcome strain (WS) and *Trypanosoma brucei brucei* ILtad 1.4 strain (IL). No difference was found in activity of ornithine decarboxylase (ODC), a key enzyme in polyamine synthesis in trypanosomes, in both strains maintained in vitro; higher ( $P < 0.05$ ) ODC values were found in IL in vivo. However, WS in vivo exhibited higher proliferation rates with higher spermidine content and decreased host survival times than IL. The in vitro proliferation and polyamine contents of WS increased with the addition of polyamine to the 1-difluoromethylornithine culture medium, but not IL. These results suggested that WS uses extracellular polyamine for proliferation. In the in vitro culture, WS was less tolerant of hydrogen peroxide (oxidative stress) than IL, and malondialdehyde levels in WS were higher than in IL. The expression of trypanothione synthetase mRNA in WS in vitro was higher than in IL. These results suggest that IL is dependent on the synthesis of polyamines for proliferation and reduction of oxidative stress, whereas WS is dependent on the uptake of extracellular polyamines. A thorough understanding of the differences in the metabolic capabilities of various trypanosomes is important for the design of more effective medical treatments.

4/7/8

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

19015652 BIOSIS NO.: 200600361047

Unnoticed spread of class 1 integrons in gram-positive clinical strains isolated in Guangzhou, China

AUTHOR: Shi Lei; Zheng Meiping; Xiao Zenghuang; %Asakura Masahiro%; Su Jianyu; Li Lin; %Yamasaki Shinji% (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, 1-1 Gakuencho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbiology and Immunology 50 (6): p463-467 2006 2006

ISSN: 0385-5600

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: A total of 46 gram-positive bacteria isolated from clinical specimens collected in China were subjected to PCR analysis with the int11-specific primers, and the int11-positive strains were further

analyzed for their resistance gene cassette. All isolates possessed the class 1 integron in their genomes and the array of gene cassettes was dhfrXII-orJF-aadA2, which is very similar to other organisms except in one isolate carrying an additional copy of the class 1 integron containing the aadA2 gene cassette. Altogether, the results indicate that the class 1 integron is widespread in gram-positive clinical strains isolated in Guangzhou, China.

4/7/9

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

18924440 BIOSIS NO.: 200600269835

Cytolethal distending toxin (CDT): Genetic diversity, structure and role in diarrheal disease

AUTHOR: %%Yamasaki Shinji%% (Reprint); %%Asakura Masahiro%%; Tsukamoto Teizo; Faruque Shah M; Deb Reema; Ramamurthy T

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Dept Vet Sci, 1-1 Gakuen Cho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Toxin Reviews 25 (1): p61-88 APR-JUN 2006 2006

ISSN: 0731-3837\_(print) 1525-6057\_(electronic)

DOCUMENT TYPE: Article; Literature Review

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: In 1987, cytolethal distending toxin (CDT) was discovered by Johnson and Lior as a new type of protein toxin produced by certain strains of *Escherichia coli*, which is different from heat-labile enterotoxin (LT) produced by enterotoxigenic *E. coli*. Although LT causes only cell elongation, CDT causes cell elongation, cell distention, irreversible cell cycle arrest, and consequently, death of the cultured mammalian cells. Recently, CDT was recognized as a new family of bacterial toxin, as a genotoxin, produced by a diverse group of gram-negative bacteria, all of which are related to mucosal infection. Although tremendous efforts have been made to study the structure and mode of action of CDT, its role in bacterial pathogenesis still remains unclear. In this review, we focus mainly on CDT produced by enteric bacteria and describe the history of CDT, their gene and protein structure, structure-function relationship, and its mode of action particularly how CDT contributes to the gastrointestinal infections.

4/7/10

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

18320405 BIOSIS NO.: 200510014905

Effects of heparin administration on *Trypanosoma brucei gambiense* infection in rats

AUTHOR: Nishimura Kazuhiko (Reprint); Shima Kensuke; %%Asakura Masahiro%%; Ohnishi Yoshihiro; %%Yamasaki Shinji%%

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Agr and Biol Sci, Div Vet Sci, 1-1, Gakuencho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: nisimura@vet.osakafu-u.ac.jp

JOURNAL: Journal of Parasitology 91 (1): p219-222 FEB 05 2005

ISSN: 0022-3395  
DOCUMENT TYPE: Article  
RECORD TYPE: Abstract  
LANGUAGE: English

ABSTRACT: We examined whether heparin administration influences in vivo trypanosome proliferation in infected rats. Administration of heparin every 8 hr via cardiac catheter inhibited growth of *Trypanosoma brucei* gambiense and prolonged survival of treated rats. Heparin administration increased lipoprotein lipase activity, high-density lipoprotein (HDL) concentration in the blood, and haptoglobin messenger RNA content of the liver. The presence of heparin in culture media did not directly affect proliferation of trypanosomes in vitro. However, the addition of plasma from infected rats treated with heparin to culture media decreased the number of trypanosomes. This effect was decreased by incubating the trypanosomes with benzyl alcohol, a known inhibitor of receptor-mediated endocytosis of lipoprotein. These data suggested that heparin administration reduced the number of trypanosomes in infected rats. Trypanosome lytic factor, a HDL and haptoglobin-related protein, protects humans and some animals from infection by *Trypanosoma brucei brucei*. In rats, increases in HDL and haptoglobin may affect the proliferation of *T. b. gambiense*.

? ds

Set	Items	Description
S1	21	(DISTENDING) AND (CAMPYLOBACTER()COLI)
S2	87	AU='YAMASAKI SHINJI'
S3	15	AU='ASAKURA MASAHIRO'
S4	10	S2 AND S3

? s s1 and s2

21	S1
87	S2
S5	3 S1 AND S2

? t s5/7/1-3

5/7/1

DIALOG(R)File 5:Biosis Previews(R)  
(c) 2008 The Thomson Corporation. All rights reserved.

0020130234 BIOSIS NO.: 200800177173

Development of a cytolethal toxin (cdt) gene-based species-specific multiplex PCR assay for the detection and identification of *Campylobacter jejuni*, *Campylobacter coli* and *Campylobacter fetus*

AUTHOR: Asakura Masahiro; Samosornsuk Worada; Hinenoya Atsushi; Misawa Naoaki; Nishimura Kazuhiko; Matsuhisa Akio; Yamasaki Shinji  
(Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Nakaku, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: FEMS Immunology and Medical Microbiology 52 (2): p260-266 MAR 2008 2008

ITEM IDENTIFIER: doi:10.1111/j.1574-695X.2007.00369.x

ISSN: 0928-8244

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: A cytolethal toxin (cdt) gene-based species-specific multiplex PCR assay for the detection of cdtA, cdtB or cdtC gene of *Campylobacter jejuni*, *Campylobacter coli* or *Campylobacter fetus*, respectively, was developed and evaluated with 76 *Campylobacter* strains belonging to seven different species and 131 other bacterial strains of eight different genera. The cdtA, cdtB or cdtC gene of *C. jejuni*, *C. coli* or *C. fetus*, respectively, could be successfully amplified using the corresponding set of primers in a highly species-specific manner. Furthermore, the specific primer set for the cdtA, cdtB or cdtC gene of a particular species could amplify the desired gene from a mixture of DNA templates of any of two or all three species. The detection limit of *C. jejuni*, *C. coli* or *C. fetus* was 10-100 CFU tube<sup>-1</sup> by the multiplex PCR assay on the basis of the presence of the cdtA, cdtB or cdtC gene. These data indicate that the cdt gene-based multiplex PCR assay may be useful for rapid and accurate detection as well as identification of *Campylobacter* strains in a species-specific manner.

5/7/2

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

0019895878 BIOSIS NO.: 200700555619

Evaluation of a cytolethal toxin (cdt) gene-based species-specific multiplex PCR assay for the identification of *Campylobacter* strains isolated from poultry in Thailand

AUTHOR: Samosornsuk Worada; Asakura Masahiro; Yoshida Emi; Taguchi Takashi; Nishimura Kazuhiko; Eampokalap Boonchuay; Phongsisay Vongsavanh; Chaicumpa Wanpen; Yamasaki Shinji (Reprint)

AUTHOR ADDRESS: Osaka Prefecture Univ, Grad Sch Life and Environm Sci, Nakaku, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbiology and Immunology 51 (9): p909-917 2007 2007

ISSN: 0385-5600

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: We have recently developed a cytolethal toxin (cdt) gene-based species-specific multiplex PCR assay for identifying *Campylobacter jejuni*, *C. coli* and *C. fetus*. In the present study, the applicability of this assay was evaluated with 34 *Campylobacter*-like organisms isolated from poultry in Thailand for species identification and was compared with other assays including API Campy, 16S rRNA gene sequence, and hippuricase (hipO) gene detection. Of the 34 strains analyzed, 20, 10 and 1 were identified as *C. jejuni*, *C. coli*, and *Arcobacter cryaerophilus*, respectively, and 3 could not be identified by API Campy. However, 16S rRNA gene analysis, showed that all 34 strains are *C. jejuni/coli*. To discriminate between these 2 species, the hipO gene, which is specifically present in *C. jejuni*, was examined by PCR and was detected in 20 strains, which were identified as *C. jejuni* by API Campy but not in the remaining 14 strains. Collective results indicated that 20 strains were *C. jejuni* whereas the 14 strains were *C. coli*. When the cdt gene-based multiplex PCR was employed, however, 19, 20 and 19 strains were identified as *C. jejuni* while 13, 14 and 13 were identified

as *C. coli* by the *cdtA*, *cdtB* and *cdtC* gene-based multiplex PCR, respectively. Pulsed-field gel electrophoresis revealed that *C. jejuni* and *C. coli* strains analyzed are genetically diverse. Taken together, these data suggest that the *cdt* gene-based multiplex PCR, particularly *cdtB* gene-based multiplex PCR, is a simple, rapid and reliable method for identifying the species of *Campylobacter* strains.

5/7/3

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019752944 BIOSIS NO.: 200700412685

Comparative analysis of cytolethal *distending* toxin (*cdt*) genes among *Campylobacter jejuni*, *C. coli* and *C. fetus* strains

AUTHOR: Asakura Masahiro; Samosornsuk Worada; Taguchi Masumi; Kobayashi Kazuhiro; Misawa Naoaki; Kusumoto Masahiro; Nishimura Kazuhiko; Matsuhisa Akio; *Yamasaki Shinji* (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Grad Sch Life and Environm Sci, Naka Ku, Gakuen Cho, Sakai, Osaka 5998531, Japan\*\*Japan

AUTHOR E-MAIL ADDRESS: shinji@vet.osakafu-u.ac.jp

JOURNAL: Microbial Pathogenesis 42 (5-6): p174-183 MAY-JUN 2007 2007

ITEM IDENTIFIER: doi:10.1016/j.micpath.2007.01.005

ISSN: 0882-4010

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The cytolethal *distending* toxin (*cdt*) gene clusters of *Campylobacter coli* strain Col-243 and *C. fetus* strain Col-187 were cloned and sequenced to understand the importance of Cdt as a virulence factor. The *cdt* genes of *C. coli* and *C. fetus* consist of three closely linked genes termed *cdtA*, *cdtB*, *cdtC* whose sizes are 774, 801, and 570 bp, and 702, 798, and 546 bp, respectively. The homologies of each subunit of *cdt* genes between *C. jejuni* and *C. coli*, *C. jejuni* and *C. fetus*, or *C. coli* and *C. fetus* are 59.6%, 40.3%, or 46.5% for *cdtA*, 70.2%, 62.4%, or 61.3% for *cdtB*, 61.3%, 52.3%, or 50.1% for *cdtC*, respectively. Colony hybridization assay revealed that the genes homologous to the *cdtABC* gene were distributed in all 27, 19, 20 strains of *C. jejuni*, *C. coli*, and *C. fetus*, respectively, isolated from patients and animals in species-specific manner. Furthermore, nucleotide sequence of the *cdt* operon, including flanking region, of 10 strains of each species indicated that though the size of the *cdtB* gene was conserved in each species, those of *cdtA* and *cdtC* genes varied particularly among *C. coli* strains. Amino acid residues demonstrated to be important for toxin activity in CdtB, corresponding to H 152, D185, D222, D258, H259 in Cj-CdtB, were also conserved in Cc-CdtB and Cf-CdtB. The *cdt* gene cluster was located in different sites among different species but in the same site of genomes of the same species. Cdt activity produced by *C. jejuni* and *C. fetus* varied among strains, however, any *C. coli* strains exhibited Cdt activity on HeLa cells. These data indicate that the *cdt* gene may have a potential for virulence factor at least in *C. jejuni* and *C. fetus*.

(C) 2007 Elsevier Ltd. All rights reserved.

? ds

Set	Items	Description
S1	21	(DISTENDING) AND (CAMPYLOBACTER())COLI)

S2 87 AU='YAMASAKI SHINJI'  
S3 15 AU='ASAKURA MASAHIRO'  
S4 10 S2 AND S3  
S5 3 S1 AND S2

? s s2 and cdta

87 S2

435 CDTA

S6 3 S2 AND CDTA

? s s6 not s5

3 S6

3 S5

S7 0 S6 NOT S5

? s s1 and cdta

21 S1

435 CDTA

S8 8 S1 AND CDTA

? s s8 not s5

8 S8

3 S5

S9 5 S8 NOT S5

? t s9/7/1-5

9/7/1

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0020123865 BIOSIS NO.: 200800170804

Optimisation of glycan and small molecule arrays for analysis of  
Campylobacter chemotaxis and adherence

AUTHOR: Asakura M (Reprint)

AUTHOR ADDRESS: Univ Osaka Prefecture, Sakai, Osaka, Japan\*\*Japan

JOURNAL: Zoonoses Public Health 54 (Suppl. 1): p99 2007 2007

CONFERENCE/MEETING: 14th International Workshop on Campylobacter,  
Helicobacter and Related Organisms Rotterdam, NETHERLANDS September 02  
-05, 2007; 20070902

ISSN: 1863-1959

DOCUMENT TYPE: Meeting; Meeting Poster

RECORD TYPE: Citation

LANGUAGE: English

9/7/2

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

0019544568 BIOSIS NO.: 200700204309

Relationships between bacterial genotypes and in vitro virulence properties  
of Campylobacter jejuni and %%%Campylobacter%%% %%%coli%%% isolated from  
turkeys

AUTHOR: Haenel I (Reprint); Borrmann E; Mueller J; Alter T

AUTHOR ADDRESS: Fed Res Inst Anim Hlth, Inst Mol Pathogenesis, Naumburger  
Str 96A, D-07743 Jena, Germany\*\*Germany

AUTHOR E-MAIL ADDRESS: ingrid.haenel@fli.bund.de

JOURNAL: Journal of Applied Microbiology 102 (2): p433-441 FEB 2007 2007

ISSN: 1364-5072

DOCUMENT TYPE: Article

RECORD TYPE: Abstract



LANGUAGE: English

ABSTRACT: Campylobacter isolates from turkeys were genotyped and characterized by their in vitro virulence properties. Relationships between bacterial genotypes and virulence properties were analysed. Isolates were analysed by pulsed-field gel electrophoresis and fla typing. The toxin production was determined on the phenotypic level using a CHO-K1 cell culture model and on the genotypic level using PCR for detection of the *cdtA*, *cdtB* and *cdtC* genes. Although the *cdtB* gene was detected from 100% of the Campylobacter jejuni and Campylobacter coli isolates we observed three different morphological pictures on the cells. Cytotoxicity was associated with cell distension or cell rounding. All four Camp. coli strains and one Camp. jejuni strain did not produce any cytotoxic changes on the cells. Adhesion, invasion and survival of Campylobacter isolates were determined in a Caco-2 cell culture model. All isolates adhered to and invaded Caco-2 cells, whereas 64.7% of the strains survived for 48 h in the cells. Seventeen Campylobacter isolates from turkeys were classified into four groups with regard to their in vitro abilities. Jackknife analysis revealed a strong association between these groups and genotype clusters. Typing methods have generally failed to identify strains with specific virulence properties. This study suggests that a relationship between subgroups of Campylobacter with common in vitro virulence characteristics and genotypes exist.

9/7/3

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rights reserved.

19060686 BIOSIS NO.: 200600406081

Adherence to and invasion of human intestinal epithelial cells by Campylobacter jejuni and Campylobacter coli isolates from retail meat products

AUTHOR: Zheng JIE; Meng JIANGHONG; Zhao SHAOHUA; Singh RUBY; Song WENXIA (Reprint)

AUTHOR ADDRESS: Univ Maryland, Dept Cell Biol and Mol Genet, College Pk, MD 20742 USA\*\*USA

AUTHOR E-MAIL ADDRESS: wenxsong@umd.edu

JOURNAL: Journal of Food Protection 69 (4): p768-774 APR 2006 2006

ISSN: 0362-028X

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The abilities of 34 Campylobacter jejuni and 9

Campylobacter coli isolates recovered from retail meats to adhere to and invade human intestinal epithelial T84 cells were examined and compared with those of a well-characterized human clinical strain, C jejuni 81-176, to better assess the pathogenic potential of these meat isolates. The meat isolates exhibited a wide range of adherence and invasion abilities; a few of the isolates adhered to and invaded T84 cells almost as well as did C jejuni 81-176. There was a significant correlation between the adherence ability and the invasion ability of the Campylobacter isolates. The presence of eight putative virulence genes in these Campylobacter isolates that are potentially responsible for adherence and invasion or that encode cytotoxic toxin

was determined using PCR. All *Campylobacter* isolates possessed *flaA*, *cadF*, *pldA*, *cdtA*, *cdtB*, and *cdtC*, and most (91%) also contained the *ciaB* gene. However, the *virBII* gene, carried by virulence plasmid pVir, was absent in almost all the *Campylobacter* isolates. Our findings indicated that *C. jejuni* and *C. coli* present in retail meat were diverse in their ability to adhere to and invade human intestinal epithelial cells and that the putative virulence genes were widespread among the *Campylobacter* isolates. Thus, despite of the presence of the putative virulence genes, only some but not all *Campylobacter* strains isolated from retail meat can effectively invade human intestinal epithelial cells in vitro.

9/7/4

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

17391700 BIOSIS NO.: 200300350419

PCR detection of seven virulence and toxin genes of *Campylobacter jejuni* and *Campylobacter coli* isolates from Danish pigs and cattle and cytolethal distending toxin production of the isolates.

AUTHOR: Bang D D (Reprint); Nielsen E Moller; Scheutz F; Pedersen K; Handberg K; Madsen M

AUTHOR ADDRESS: Department of Poultry, Fish and Fur Animals, Danish Veterinary Institute, Hangevej 2, DK-8200, Aarhus N, Denmark\*\*Denmark

AUTHOR E-MAIL ADDRESS: ddb@vetinst.dk

JOURNAL: Journal of Applied Microbiology 94 (6): p1003-1014 2003 2003

MEDIUM: print

ISSN: 1364-5072

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: Aims: To study the prevalence of seven virulence and toxin genes, and cytolethal distending toxin (CDT) production of *Campylobacter jejuni* and *C. coli* isolates from Danish pigs and cattle. Methods and Results: The presence of the *cadF*, *ceuE*, *virB11*, *flaA*, *cdtA*, *cdtB*, *cdtC* and the *cdt* gene cluster among 40 *C. jejuni* and *C. coli* isolates was detected by polymerase chain reaction. The CDT production of the isolates was determined on Vero, colon 205 and chicken embryo cells. The *cadF*, *flaA*, *ceuE* and *cdtB* genes were detected from 100% of the isolates. The *cdtA* and *cdtC* genes were found in 95.0 and 90.0% of the isolates, respectively. The *cdt* gene cluster was detected in 82.5% isolates. Only 7.5% of the isolates were positive for *virB11*. Ninety-five per cent of the isolates produced CDT in Vero and colon 205 cell assays, and 90% of the isolates produced CDT in chicken embryo cell assays. Conclusions: High prevalence of the *cadF*, *ceuE*, *flaA* and *cdtB* genes was found. Data of the prevalence of *cdt* genes was consistent with the CDT titres produced by the isolates. *Campylobacter coli* from pigs produced high CDT titres. Significance and Impact of the Study: The high prevalence of seven virulence and toxin genes demonstrated that these putative pathogenic determinants are widespread among *Campylobacter* isolates from pigs and cattle. *Campylobacter coli* isolates from pigs produced much higher CDT titres compared with *C. coli* isolates from other sources suggesting that *C. coli* may be particularly adapted to or associated with this species.

9/7/5

DIALOG(R)File 5:Biosis Previews(R)

(c) 2008 The Thomson Corporation. All rts. reserv.

13406673 BIOSIS NO.: 199699040733

Prevalence of cytolethal ~~distending~~ toxin production in *Campylobacter jejuni* and relatedness of *Campylobacter* sp. *cdtB* genes

AUTHOR: Pickett Carol L (Reprint); Pesci Evertt C; Cottle Daniel L; Russell Gina; Erdem Aysegul Nalca; Zeytn Hasan

AUTHOR ADDRESS: Dep. Microbiol. Immunol., Univ. Kentucky, Chandler Med.

Cent., 800 Rose Street, Lexington, KY 40536-0084, USA\*\*USA

JOURNAL: Infection and Immunity 64 (6): p2070-2078 1996 1996

ISSN: 0019-9567

DOCUMENT TYPE: Article

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: *Campylobacter jejuni* produces a toxin called cytolethal ~~distending~~ toxin (CDT). The genes encoding this toxin in *C. jejuni* 81-176 were cloned and sequenced. The nucleotide sequence of the genes revealed that there are three genes, ~~cdtA~~, *cdtB*, and *cdtC*, encoding proteins with predicted sizes of 30,116, 28,989, and 21,157 Da, respectively. All three proteins were found to be related to the *Escherichia coli* CDT proteins, yet the amino acid sequences have diverged significantly. All three genes were required for toxic activity in a HeLa cell assay. HeLa cell assays of a variety of *C. jejuni* and *C. coli* strains suggested that most *C. jejuni* strains produce significantly higher CDT titers than do *C. coli* strains. Southern hybridization experiments demonstrated that the *cdtB* gene is present on a 6.0-kb *Cla*I fragment in all but one of the *C. jejuni* strains tested; the *cdtB* gene was on a 6.9-kb *Cla*I fragment in one strain. The *C. jejuni* 81-176 *cdtB* probe hybridized weakly to DNAs from *C. coli* strains. The *C. jejuni* 81-176 *cdtB* probe did not hybridize to DNAs from representative *C. fetus*, *C. lari*, *C. "upsaliensis"*, and *C. hyointestinalis* strains, although the HeLa cell assay indicated that these strains make CDT. PCR experiments indicated the probable presence of *cdtB* sequences in all of these *Campylobacter* species.

? ds

Set	Items	Description
S1	21	(DISTENDING) AND (CAMPYLOBACTER())COLI)
S2	87	AU='YAMASAKI SHINJI'
S3	15	AU='ASAKURA MASAHIRO'
S4	10	S2 AND S3
S5	3	S1 AND S2
S6	3	S2 AND CDTA
S7	0	S6 NOT S5
S8	8	S1 AND CDTA
S9	5	S8 NOT S5

? log y

04sep08 09:34:15 User217744 Session D1155.3

\$15.51 2.506 DialUnits File5

\$95.16 39 Type(s) in Format 7

\$95.16 39 Types

\$110.67 Estimated cost File5

\$2.40 TELNET

\$113.07 Estimated cost this search  
\$113.09 Estimated total session cost 2.886 DialUnits  
Logoff: level 05.22.00 D 09:34:15